

SUSSEX SHOULDER & ELBOW

06

Acromioclavicular Joint Dislocation

What is Acromioclavicular joint (ACJ) dislocation?

The ACJ is the small fibrous joint between the end of your clavicle (collar bone) and the acromion (a protruberance of your wing-blade). You can usually feel your ACJ as the small bump on the top of your shoulder. If you fall heavily on the side of your shoulder, such as when falling off your bike or during rugby, the impact can compress the acromion beneath the end of the clavicle forcing it upwards. Depending on the force of the impact the joint will either be sprained or will actually dislocate. The injury is typically graded into a Type I injury (simple compression of the joint), Type II where the acromio-clavicular ligament is damaged and there is slight widening of the joint and Type III where all the ligaments stabilising the joint are damaged and the end of the clavicle displaces 100%. There are also other less common types of injury.

How is ACJ dislocation diagnosed?

The injury is diagnosed from the history of injury and examination of the shoulder. A thorough examination is very important to exclude other injuries to the shoulder. There is often an obvious bump on the top of the shoulder which is very tender. Movement of the shoulder is limited by pain. An x-ray will demonstrate, in most cases, the injury to the joint and will exclude a fracture of the end of the clavicle. An MRI scan is seldom indicated.

What are the treatment options?

For most Type I and II injuries the shoulder is rested in a sling for two to three weeks whilst the joint is slowly mobilised as pain allows. Ice and painkillers are helpful. You will then be referred for physiotherapy to help restore movement and strengthen the muscles around the shoulder. It can take several months to recover good function in your shoulder. Occasionally a steroid injection into the injured joint is required to settle persistent discomfort. About 5-10% of patients will have ongoing problems such as painful clicking a year after injury and, in those patients, we usually recommend key hole excision of the end of the clavicle and clearance of the damaged cartilage. Most type III injuries can also be managed without surgery initially. In our experience if you have not made quite good progress by three or four months post injury then we would recommend surgical reconstruction of the ACJ. For more severe grades of injury we recommend early reconstruction.

What does surgery involve?

For persistent pain after a Type I or Type II injury we usually recommend key-hole surgery to excise 5mm off the end of the collar bone and clear the scar tissue. You will be able to move your arm straight after surgery but it can take up to three months for the pain to settle. For reconstruction of Type III or more severe injuries we use a ligament graft which may be synthetic or tissue based. This graft loops around the coracoid (a protuberance of your wing-blade), passes behind your collar bone and is fixed in place on the top of your collar bone with a screw once the collar bone has been pulled down into place.

Cameron Hatrick

MA (Cantab), MB, BChir, FRCS (Tr&Orth)
Consultant Orthopaedic Surgeon

Joideep Phadnis

FRCS (Tr&Orth), Dip Sports Med, MRCS, MBChB
Consultant Orthopaedic Surgeon

These notes are intended as a guide and some of the details may vary depending on your individual circumstance and at the discretion of your surgeon.
© Sussex Shoulder

Acromioclavicular Joint Dislocation

What can I expect after surgery?

You will wake up from surgery with your arm in a sling which you should wear for the next three weeks. Your arm will feel numb and 'heavy' whilst the nerve block is working during the first night. The shoulder will become a bit sore after that but you will be provided with painkillers which you should take regularly for the first few days. For the first three weeks you will exercise your elbow and shoulder carefully, taking care not to lift it above the horizontal. You should avoid lifting with your arm for the first 6 weeks. You will be provided with a 'rehabilitation' sheet showing you the appropriate exercises or you can download the instructions from the Exercises and Rehabilitation section of this website.

Most patients will have recovered quite good movement by 6 weeks post surgery, will be able to return to driving by 4-6 weeks, will be able to return to light manual work by 6 weeks and heavier duties, including sports, by 12 weeks. By 3 months 80% patients can expect to have had a good or excellent outcome.

Cameron Hatrick

NHS Patients

Brighton NHS
Tel: 01273 696955 ext 4258, Tara Kirk
Lewes NHS
Tel: 01273 474153 ask for Glynis Mockett

Private Patients

Direct Tel: 01273 828095 Angela Crosskey
Montefiore Hospital Hove
Tel: 01273 828120

Joideep Phadnis

NHS Patients

Brighton NHS
Tel: 01273 696955 ext 8258, Shirley Boyack

Private Patients

Direct Tel: 01273 828099, Allison Litynski
Montefiore Hospital Hove
Tel: 01273 828120

We value your feedback

myclinicaloutcomes.co.uk

Don't forget to register on myclinicaloutcomes.co.uk before your surgery so that you can track the progress of your shoulder.

iwantgreatcare.org

You are also welcome to provide feedback on iwantgreatcare.org so that we can continue to improve the service we provide.