

SUSSEX SHOULDER & ELBOW

01

Arthroscopic Subacromial Decompression and/or ACJ Excision

What happens before surgery?

Once a decision has been made to proceed to surgery you will be contacted by staff at the hospital where you are going to have your procedure for a pre-operative assessment. This will either take place on the telephone or at the hospital if you have any medical problems that might require a more detailed assessment.

What happens on the day of surgery?

You will be admitted to the ward or the day surgery unit, usually on the morning of your surgery. You will be seen by the anaesthetist and by your surgeon. They will take you through the details of the anaesthetic, which usually includes a nerve block to help your pain, and through the details of the operation. You will be asked to sign a consent form and to complete a pre-operative shoulder score so that we can track your progress following surgery.

What does the surgery involve?

The operation involves the trimming of bone from the front of the acromion bone, to increase space for the tendons. If the acromio-clavicular joint (ACJ) is also causing pain then a few mm of bone will also be shaved from the end of the clavicle to decompress the ACJ. Occasionally other damage is found within the joint that requires repair.

What happens after the operation?

You will wake up in the recovery unit where a nurse will be looking after you. Your arm will be in a sling and will usually feel very heavy and numb as a result of the nerve block. The shoulder is usually quite comfortable but you will be given painkillers if you are in any pain. Once you have recovered from your anaesthetic you will be transferred back to the ward or to the 'discharge' area.

A physiotherapist will then see you and take you through how to apply and remove your sling safely and the initial exercises you should perform. Once you are comfortable you will be sent home, usually on the day of surgery, with painkillers and instructions about any stitches you have. You will need to make an appointment with your practice nurse to have the stitches removed or trimmed after 8 days. You will be sent an appointment to be seen in the clinic 2-3 weeks after your operation. A physiotherapy appointment, if required, will be arranged by the hospital.

Cameron Hatrick

MA (Cantab), MB, BChir, FRCS (Tr&Orth)
Consultant Orthopaedic Surgeon

Joideep Phadnis

FRCS (Tr&Orth), Dip Sports Med, MRCS, MBChB
Consultant Orthopaedic Surgeon

These notes are intended as a guide and some of the details may vary depending on your individual circumstance and at the discretion of your surgeon.
© Sussex Shoulder

Arthroscopic Subacromial Decompression and/or ACJ Excision

What exercises should I perform after the operation?

1st 2 weeks:

- Wear your sling for a few days for comfort, but you should aim to discard it by the end of the first week.
- Remove the sling 3-5 times each day to perform the following exercises:
- Active finger, wrist and elbow movements
- Shoulder pendulum exercises for the first day or two followed by:
- Full active assisted and active shoulder exercises as pain allows
- Scapular setting exercises

2 weeks onwards:

- Progress active range of movement
- Gradually increase rotator cuff strengthening exercises

What is the usual recovery?

- 2-3 weeks: driving
- 3 weeks: approximately 75% of movement but with pain
- 6 weeks: nearly full range of movement
- 8 weeks: golf
- 12 weeks: swimming, racquet sports
- 3-6 months: full recovery. Pain, particularly at night, can persist

Return to work: depends on occupation

- Home based sedentary work (e.g computer) from one week
- Sedentary work in an office from one to two weeks
- Light manual work from three to four weeks
- Heavy manual work from six-eight weeks

Are there any complications of surgery?

Fortunately complications after shoulder surgery are uncommon.

They include:

- Infection (< 1 in 1000)
- Stiffness. Mild stiffness is quite common but occasionally a full frozen shoulder can develop (5%) which will prolong your recovery by a few months.
- Pain. This is common for the first few weeks after surgery but steadily settles. Some patients will have ongoing discomfort at 3 months that requires a post-operative steroid injection to settle the inflammation.
- Fracture. If too much bone is shaved from the acromion it may fracture and require fixation. This is very rare.
- Failure to improve. 80% of patients will make a good or excellent recovery. 15% will have some ongoing discomfort but will be satisfied with their outcome. About 5% of patients will have ongoing problems, such as pain and stiffness, and some of these require further surgery.

Cameron Hatrick

NHS Patients

Brighton NHS
Tel: 01273 696955 ext 4258, Tara Kirk
Lewes NHS
Tel: 01273 474153 ask for Glynis Mockett

Private Patients

Direct Tel: 01273 828095 Angela Crosskey
Montefiore Hospital Hove
Tel: 01273 828120

Joideep Phadnis

NHS Patients

Brighton NHS
Tel: 01273 696955 ext 8258, Shirley Boyack

Private Patients

Direct Tel: 01273 828099, Allison Litynski
Montefiore Hospital Hove
Tel: 01273 828120

We value your feedback

myclinicaloutcomes.co.uk

Don't forget to register on myclinicaloutcomes.co.uk before your surgery so that you can track the progress of your shoulder.

iwantgreatcare.org

You are also welcome to provide feedback on iwantgreatcare.org so that we can continue to improve the service we provide.